

RESEARCH

Open Access



Observational study of lenalidomide in patients with mantle cell lymphoma who relapsed/progressed after or were refractory/intolerant to ibrutinib (MCL-004)

Michael Wang^{1*}, Stephen J. Schuster², Tycel Phillips³, Izidore S. Lossos⁴, Andre Goy⁵, Simon Rule⁶, Mehdi Hamadani⁷, Nilanjan Ghosh⁸, Craig B. Reeder⁹, Evelyn Barnett¹⁰, Marie-Laure Casadebaig Bravo¹¹ and Peter Martin¹²

Abstract

Background: The observational MCL-004 study evaluated outcomes in patients with relapsed/refractory mantle cell lymphoma who received lenalidomide-based therapy after ibrutinib failure or intolerance.

Methods: The primary endpoint was investigator-assessed overall response rate based on the 2007 International Working Group criteria.

Results: Of 58 enrolled patients (median age, 71 years; range, 50–89), 13 received lenalidomide monotherapy, 11 lenalidomide plus rituximab, and 34 lenalidomide plus other treatment. Most patients (88%) had received ≥ 3 prior therapies (median 4; range, 1–13). Median time from last dose of ibrutinib to the start of lenalidomide was 1.3 weeks (range, 0.1–21.7); 45% of patients had partial responses or better to prior ibrutinib. Primary reasons for ibrutinib discontinuation were lack of efficacy (88%) and ibrutinib toxicity (9%). After a median of two cycles (range, 0–11) of lenalidomide-based treatment, 17 patients responded (8 complete responses, 9 partial responses), for a 29% overall response rate (95% confidence interval, 18–43%) and a median duration of response of 20 weeks (95% confidence interval, 2.9 to not available). Overall response rate to lenalidomide-based therapy was similar for patients with relapsed/progressive disease after previous response to ibrutinib (i.e., \geq PR) versus ibrutinib-refractory (i.e., \leq SD) patients (30 versus 32%, respectively). The most common all-grade treatment-emergent adverse events after lenalidomide-containing therapy ($n = 58$) were fatigue (38%) and cough, dizziness, dyspnea, nausea, and peripheral edema (19% each). At data cutoff, 28 patients have died, primarily due to mantle cell lymphoma.

Conclusion: Lenalidomide-based treatment showed clinical activity, with no unexpected toxicities, in patients with relapsed/refractory mantle cell lymphoma who previously failed ibrutinib therapy.

Trial registration: Clinicaltrials.gov identifier NCT02341781. Date of registration: January 14, 2015

Keywords: Ibrutinib failure, Lenalidomide, Mantle cell lymphoma

* Correspondence: miwang@mdanderson.org

¹Department of Lymphoma/Myeloma, University of Texas MD Anderson Cancer Center, 1515 Holcombe Boulevard, Houston, TX, USA
Full list of author information is available at the end of the article

Background

Mantle cell lymphoma (MCL) accounts for 3 to 6% of non-Hodgkin lymphomas and is generally characterized by cyclin D1 overexpression and, more recently, by SOX11 expression [1–3]. MCL is generally considered incurable with standard chemoimmunotherapy and approved targeted agents [4]. Although multiple molecular-based therapies have improved outcomes for patients with relapsed/refractory MCL, there is no established standard-of-care [5, 6]. As summarized in a recent review, various chemoimmunotherapy regimens tested in small clinical trials in this setting have achieved high overall response rates (ORR) ranging from 58 to 93%, but progression-free survival (PFS) has been limited to < 2 years [6], with reported overall survival (OS) as < 3 years [7–9].

Bortezomib, lenalidomide, and ibrutinib have received US Food and Drug Administration (FDA) approval for the treatment of relapsed/refractory MCL [10–12], and lenalidomide, ibrutinib, and temsirolimus are registered for this indication in the European Union [10, 13, 14]. Monotherapy activities with these targeted agents in phase II studies report ORRs ranging from 22 to 68%, complete response (CR) rates ranging from 2 to 21%, and median duration of response (DOR) ranging from 9.2 to 19.6 months [6]. In a randomized study comparing two targeted agents in patients with relapsed/refractory MCL, ibrutinib significantly reduced the risk of progressive disease (PD) or death compared with temsirolimus (hazard ratio [HR] 0.43; 95% confidence interval [CI], 0.32–0.58; $P < 0.0001$) [15]. After a median follow-up of 20 months, ibrutinib demonstrated an improved median PFS (14.6 versus 6.2 months; $P < 0.0001$), 2-year PFS (41 versus 7%; P value not reported), ORR (72 versus 40%; $P < 0.0001$), and CR rate (19 versus 1%; P value not reported) compared with temsirolimus.

Although these treatments have shown significant antitumor activity and are commonly used, primary and acquired resistance, intolerance, and drug-related toxicities are significant limitations of these treatment approaches. With ibrutinib in particular, recent studies have shown that MCL patients with primary or acquired resistance have poor clinical outcomes. A retrospective review of 31 patients with MCL who had PD following discontinuation of ibrutinib and received salvage chemoimmunotherapy showed an ORR of 32% with the first salvage regimen and an estimated 22% 1-year OS (median 8.4 months) [16]. In another retrospective analysis, 114 heavily pretreated patients with MCL developed PD, while on ibrutinib (for a median treatment duration of 4.7 months) had a median OS of 2.9 months after discontinuing ibrutinib [17].

The oral immunomodulatory drug (IMiD[®]) lenalidomide has demonstrated antitumor activity in preclinical studies of MCL, both as monotherapy and in combination with

rituximab [18–21]. In clinical trials in patients with relapsed/refractory MCL and other non-Hodgkin lymphomas, lenalidomide demonstrated activity when used as a monotherapy [22–28] and in combination with rituximab (R²) [29, 30].

The objective of this retrospective, observational, multi-center MCL-004 study (NCT02341781) was to evaluate the clinical effectiveness and safety of lenalidomide used as monotherapy and in combination regimens to treat patients with MCL who had relapsed/progressed to an ibrutinib-containing treatment (i.e., had an initial response of PR or better) or who were refractory to (i.e., best response of SD or worse) or unable to tolerate ibrutinib.

Methods

Patients

Harmonization E6 requirements (Good Clinical Practice) and ethical principles per the Declaration of Helsinki were followed. All aspects of the study were reviewed with the study investigators and staff; accuracy was confirmed through source data verification.

Inclusion criteria were age ≥ 18 years; MCL verified by investigator review of a pathology report; at least one dose (cycle 1, day 1) of ibrutinib (monotherapy or combination); and ibrutinib failure defined as relapse (CR followed by relapse at any time), PD (PR followed by PD at any time), refractory (PD, or stable disease [SD] followed by PD, while on ibrutinib), and/or intolerance (discontinuation of ibrutinib for reasons other than PD). Lenalidomide was not required to immediately follow ibrutinib.

Study design

After identifying MCL patients treated with or intending to take lenalidomide following ibrutinib failure, an informed consent document was completed by the patient (family member/legal representative if patient was deceased), or a waiver was granted from the Institutional Review Board or Ethics Committee (IRB/EC) if consent was deemed not necessary for data collection. Patients were then enrolled into the clinical database, and data were extracted from medical charts including demographic information, relevant medical history, baseline disease characteristics, date of initial MCL diagnosis with pathology report, prior therapies (including treatment dates and best response), ibrutinib and lenalidomide treatment dates and outcome, copy of imaging reports, date of last follow-up/disease status, documentation of adverse events (AEs), and date/cause of death. Patients were enrolled after meeting eligibility criteria. Non-retrospective data may have been collected when lenalidomide was ongoing at study entry.

The primary endpoint was ORR defined as achievement of CR or PR per 2007 International Working Group (IWG) response criteria [31]. When initial

assessments used IWG 1999 criteria (i.e., unconfirmed CR [32]), the corresponding response per IWG 2007 was changed to PR. Patients without a response evaluation or had an unknown response were considered non-responders. The secondary endpoint was DOR (time from initial response to lenalidomide-based therapy of \geq PR to relapse/PD/death, whichever occurred first). Responding patients without PD/death at analysis were censored at the last assessment date.

Response and safety assessments

Time-to-event data were estimated using the Kaplan-Meier method [33]. Planned analyses were conducted for MCL subgroups of refractory (best response to ibrutinib of SD or worse), relapsed/PD (initial response to ibrutinib of \geq PR followed by PD), and those unable to tolerate ibrutinib (any reason other than lack of efficacy).

Available records of treatment-emergent AEs (TEAEs) with an onset date after lenalidomide initiation through 28 days after the last lenalidomide dose, regardless of causality, were analyzed in the safety population. AEs were classified according to the National Cancer Institute (NCI) Common Terminology Criteria for Adverse Events (CTCAE) version 4.03.

Statistical analysis

All efficacy evaluations were conducted in the eligible patients. Patients were grouped by first type of lenalidomide treatment received: single agent, in combination with rituximab, or in combination with other agents. The response rate probability was estimated using the proportion of responding patients with an exact two-sided 95% CI; a sample size of 30 patients would allow a two-sided 95% CI (lower boundary of 10%) for an expected proportion of 25%.

Results

Patient characteristics

MCL patients from March 1, 2009, to April 12, 2016 who were treated with lenalidomide following ibrutinib therapy were enrolled. The data cutoff for all patients was November 1, 2016. The study enrolled 58 patients at a total of 11 study sites, including 10 sites in the USA and one site in England (Additional file 1: Table S1). Seven patients signed informed consent forms (one patient signed consent prior to initiating lenalidomide treatment), and 51 patients had IRB/EC waivers. Thirteen patients were treated with lenalidomide monotherapy, 11 with lenalidomide plus rituximab, and 34 with other lenalidomide combinations (Additional file 1: Table S2). Two patients initially identified for analysis were excluded from this observational cohort because they did not meet all eligibility criteria (one patient treated with lenalidomide plus rituximab had not

relapsed while on ibrutinib and one patient was not treated with lenalidomide); these two patients are not included in the overall enrolled set of 58 patients.

Patients had a median age of 71 years (range, 50–89), and 71% were aged \geq 65 years (Table 1). Forty-eight percent of patients had an Eastern Cooperative Oncology Group performance status of 0–1, 29% had high tumor burden, and 14% had bulky disease (\geq 7 cm). The Mantle Cell International Prognostic Index (MIPI) score could not be derived for most patients due to a lack of the required data to complete appropriate calculations for 30 patients (i.e., 52% missing data for MIPI; Ki-67 data were not collected).

Patients had received a median of four prior lines of systemic anti-lymphoma therapy (range, 1–13), 88% had three or more prior therapies, and 79% had received ibrutinib as monotherapy (Table 2). Most patients (60%)

Table 1 Patient characteristics at study entry

Characteristic	L (n = 13)		L + R (n = 11)		L + other (n = 34)		Overall (N = 58)	
	No.	%	No.	%	No.	%	No.	%
Median age, years (range)	67 (54–83)		70 (58–84)		71 (50–89)		71 (50–89)	
\geq 65	6	46	9	82	26	76	41	71
Sex								
Male	11	85	8	73	25	74	44	76
Female	2	15	3	27	9	26	14	24
ECOG PS								
0–1	7	54	5	45	16	47	28	48
2–4	3	23	1	9	4	12	8	14
Missing	3	23	5	45	14	41	22	38
Tumor burden ^a								
High	4	31	1	9	12	35	17	29
Low	1	8	5	45	13	38	19	33
Missing	8	62	5	45	9	26	22	38
Bulky disease ^b								
Yes	2	15	0	0	6	18	8	14
No	2	15	6	55	17	50	25	43
Missing	9	69	5	45	11	32	25	43
Time from diagnosis to first lenalidomide dose, months								
Median	58		47		46		49	
Range	15–144		6–105		4–214		4–214	
Time from end of last prior antilymphoma therapy to first dose of lenalidomide, weeks								
Median	0.7		0.3		0.7		0.7	
Range	0.1–3.5		0.1–21.7		0.1–12.6		0.1–21.7	

ECOG PS Eastern Cooperative Oncology Group performance status, L lenalidomide, L + R lenalidomide plus rituximab

^aHigh tumor burden is defined as at least one lesion \geq 5 cm in diameter or three lesions \geq 3 cm in diameter [22]

^bBulky disease is defined as at least one lesion \geq 7 cm in the longest diameter²²

Table 2 Treatment history of enrolled patients

	L (n = 13)		L + R (n = 11)		L + other (n = 34)		Overall (N = 58)	
	No.	%	No.	%	No.	%	No.	%
No. of prior antilymphoma treatment regimens								
Median	4		3		4		4	
Range	3–7		2–8		1–13		1–13	
No. of prior antilymphoma therapies								
1	0	0	0	0	1	3	1	2
2	0	0	4	36	2	6	6	10
3	5	38	3	27	10	29	18	31
≥ 4	8	62	4	36	21	62	33	57
Missing	0	0	0	0	0	0	0	0
Type of ibrutinib treatment								
Combination regimen	1	8	1	9	10	29	12	21
Monotherapy	12	92	10	91	24	71	46	79
Ibrutinib status at study inclusion								
Relapse/PD	6	46	2	18	15	44	23	40
Refractory	2	15	8	73	15	44	25	43
Intolerant	3	23	0	0	3	9	6	10
Missing	2	15	1	9	1	3	4	7
Duration of ibrutinib treatment, months								
Median	4.8		3.9		4.3		4.3	
Range	1.2–13.9		2.0–16.6		0.5–47.6		0.5–47.6	
Best response on ibrutinib								
CR	2	15	0	0	6	18	8	14
PR	5	38	2	18	11	32	18	31
SD	0	0	1	9	0	0	1	2
Relapse/PD	5	38	8	73	15	44	28	48
Unknown	1	8	0	0	2	6	3	5
Primary reason for ibrutinib discontinuation								
Lack of efficacy	9	69	11	100	31	91	51	88
Toxicity to ibrutinib	3	23	0	0	2	6	5	9
Toxicity attribution unknown	0	0	0	0	1	3	1	2
Completed ibrutinib treatment	1	8	0	0	0	0	1	2
Time from end of last dose of ibrutinib to first dose of lenalidomide, weeks ^a								
Median	1.4		0.4		1.3		1.3	
Range	0.1–7.4		0.1–21.7		0.1–16.8		0.1–21.7	

CR complete response, L lenalidomide, L + R lenalidomide plus rituximab, PD progressive disease, PR partial response, SD stable disease

^aTime from last dose of ibrutinib to first dose of lenalidomide (weeks) is calculated as (lenalidomide first dose date – end date of ibrutinib + 1)/7

had lenalidomide-containing therapy as their next line of therapy, and 40% patients had ≥ 1 line(s) of other therapy preceding the lenalidomide regimen. Median duration of ibrutinib treatment was 4.3 months (range, 0.5–47.6).

Eighty-eight percent of patients discontinued ibrutinib treatment for one or more reason, due to relapse/PD (n = 27) and/or refractoriness (n = 25), six patients discontinued due to toxicity, and one patient completed ibrutinib as planned but had relapsed/PD at the end of ibrutinib treatment. Besides ibrutinib, the most common previous systemic therapies were rituximab (97%), cyclophosphamide (84%), glucocorticoids (78%), vincristine (78%), doxorubicin (72%), bendamustine (57%), and cytarabine (52%) (Additional file 1: Table S3; note that multiple treatment names could be used to collect this information). The median time from last dose of ibrutinib to first dose of lenalidomide was 1.3 weeks (range, 0.1–21.7) (Table 3).

Efficacy

Among the 58 patients, the median duration of treatment was 8.4 weeks for single-agent lenalidomide and 7.4 weeks for lenalidomide-containing combination therapy (Table 4). Eight patients achieved a CR and nine achieved a PR with lenalidomide-based therapy, for an ORR of 29% (95% CI, 18–43%; Table 3), which exceeded the predefined lower boundary of the 95% confidence threshold of 10% ORR. Seven of the eight patients with CR had CT ± PET/CT assessments. Two of the 13 patients (15%) who had single-agent lenalidomide (fourth line of therapy for both) reported a best response of relapse/PD to ibrutinib; 3/13 (23%) patients on single-agent lenalidomide had unknown response status with 8/13 (62%) reporting relapse/PD.

The median DOR for responders was 20 weeks (95% CI, 2.9 to not reached); of the 17 responders, 14 (82%; 7

Table 3 Efficacy outcomes with lenalidomide in patients with MCL after ibrutinib failure or intolerance

Outcome	L (n = 13)		L + R (n = 11)		L + other ^a (n = 34)		Overall (N = 58)	
	No.	%	No.	%	No.	%	No.	%
Best response by investigator’s assessment								
ORR	2	15	3	27	12	35	17	29
95% CI	2–45%		6–61%		20–54%		18–43%	
CR	0	0	1	9	7	21	8	14
PR	2	15	2	18	5	15	9	15
SD	0	0	1	9	3	9	4	7
Relapse/PD	8	62	3	27	16	47	27	47
Unknown	3	23	2	18	3	9	8	14
Missing	0	0	2	18	0	0	2	3
Duration of response, weeks								
KM median	3		20		NA		20	
95% CI	NA to NA		NA to NA		16.4 to NA		2.9 to NA	

CI confidence interval, CR complete response, KM Kaplan-Meier, L lenalidomide, L + R lenalidomide plus rituximab, MCL mantle cell lymphoma, NA not applicable, PD progressive disease, PR partial response, SD stable disease

^aAdditional file 1: Table S2 lists the other treatments

Table 4 Lenalidomide treatment exposure (safety population)

	L (n = 13)	L + R (n = 11)	L + other (n = 34)	Overall (N = 58)
Lenalidomide treatment duration, weeks				
Median	8.4	14.0	7.0	8.4
Range	0.4 to 30.0	0.9 to 37.9	1.1 to 77.9	0.4 to 77.9
Number of lenalidomide cycles				
Median	2.0	2.0	1.0	2.0
Range	1.0 to 7.0	1.0 to 9.0	0.0 to 11.0	0.0 to 11.0
Duration of other therapy combined with lenalidomide, weeks				
Median	NA	8.3	7.2	7.4
Range	NA	0.1 to 35.9	0.7 to 77.7	0.1 to 77.7

L lenalidomide, L + R lenalidomide plus rituximab, NA not applicable


CR and 7 PR) were censored from the DOR analysis due to lack of follow-up data on PD or death. At the last available assessment of the 14 censored patients, three were ongoing, three had completed lenalidomide treatment as planned, and eight patients discontinued lenalidomide treatment early (withdrew consent [$n = 1$], patient decision [$n = 1$], enrolled in a clinical trial for oral treatment [$n = 1$], started other lines of treatment [$n = 3$; because of lung cancer, physician's decision, or bone marrow transplant], and toxicity [$n = 2$]). One of the censored patients who had a first response of PR and best response of CR had the last censored DOR at 25 weeks before stopping therapy. For the three uncensored patients, two had a best response of PR and one had CR, with an estimated DOR of 2.9, 19.7, and 16.4 weeks, respectively. Univariate analysis showed a median DOR of 16 weeks (95% CI, 2.9–19.7) in the three uncensored patients (14 patient responders were censored; total of 17 responders).

Response by subgroup analysis

Patients with MCL refractory to ibrutinib versus those who relapsed/progressed on or following ibrutinib had similar ORRs of 32 versus 30%, respectively (Fig. 1); however, the CR rates were not similar (8 versus 22%). The median DOR was 20 weeks (CI 95%, 2.9–20) for the ibrutinib-refractory group and not available for the relapsed/PD group. There was one PR (17%) among the six patients who were ibrutinib-intolerant; all six patients were treated with lenalidomide within 6 months of stopping ibrutinib therapy. Of the 48 patients who tolerated ibrutinib therapy, seven had CRs and eight had PRs, a 31% ORR, and the median DOR was 20 weeks.

Safety

Overall, patients received a median of two cycles (range, 0–11) of lenalidomide-based treatment. Most patients received lenalidomide 10–25 mg/day on days 1–21 of each 28-day cycle. As of the cutoff date of November 1,



2016, 54 patients had discontinued lenalidomide-based therapy and four patients continue to receive lenalidomide (three censored for efficacy analyses), one in combination with weekly bortezomib/dexamethasone/rituximab, two in combination with weekly rituximab, and one in combination with weekly obinutuzumab. The primary reasons for lenalidomide treatment discontinuation were lack of efficacy ($n = 27$); toxicity ($n = 10$); other reasons ($n = 9$), such as initiation of another therapy (e.g., based on physician or patient choice) or trial (also an oral therapy), undergoing stem cell transplantation, or primary clinician/patient decision to stop therapy; completion of lenalidomide treatment ($n = 5$); and missing data ($n = 3$).

Of the 58 patients analyzed for safety, 48 (83%) had one or more TEAE during lenalidomide treatment. Twenty (34%) patients had at least one serious TEAE (lenalidomide alone 23%; lenalidomide + rituximab 36%; lenalidomide + others 38%). The most frequently reported serious TEAEs of any grade were febrile neutropenia ($n = 4$; 7%), hypotension (7%), deep vein thrombosis (DVT) ($n = 3$; 5%), pneumonia (5%), pancytopenia (5%), fall (5%), acute kidney injury (5%), dyspnea ($n = 2$; 3%), sepsis (3%), and respiratory failure (3%). Overall, nine (16%) patients had at least one TEAE leading to dose discontinuation (lenalidomide alone 8%; lenalidomide + rituximab 18%; lenalidomide + others 18%). These TEAEs included pancytopenia, thrombocytopenia, and rash, each experienced by two patients (3%), and anemia, febrile neutropenia, neutropenia, sepsis, fall, squamous cell lung carcinoma, dyspnea, pleural effusion, and orthostatic hypotension, each experienced by one patient (2%). The most common all-grade TEAEs were fatigue, cough, dizziness, dyspnea, nausea, peripheral edema,

anemia, rash, thrombocytopenia, and neutropenia (Table 5).

As of the cutoff date, 28 (48%) patients had died, 12 (21%) during treatment with lenalidomide, and 15 (26%) during follow-up (one unknown). Overall, 20 (34%) patients died from malignant disease (i.e., MCL) or its complications, five from unknown causes (not assessable or insufficient data), one reported another cause of end-stage renal disease, and two due to AEs. Of the two patients who died due to AEs, the first patient included a 68-year-old man in the lenalidomide-alone group who died during treatment (83 days after the first lenalidomide dose). This patient had a PR 2 months after lenalidomide initiation but died due to a pulmonary embolism, suspected to be related to lenalidomide therapy, as well as had incidences of other grade 5 AEs (DVT and cardiac arrest). Although this patient was receiving aspirin, therapy was stopped during study admission. For most patients, it is not known if the

patients received antithrombotic treatment, since concomitant treatments were not part of the collected data. The second patient who died due to an AE was a 71-year-old man who received one treatment cycle of lenalidomide in combination with ibrutinib, rituximab, bortezomib, and dexamethasone. This second patient died while on study treatment (25 days after the first dose of lenalidomide) because of progression of MCL (which included acute kidney injury, lactic acidosis, respiratory failure, and hypotension).

Discussion

This multicenter observational study examined outcomes with lenalidomide treatment in patients with MCL who had relapsed or progressed after or during ibrutinib therapy or were intolerant to ibrutinib. Most patients had received three or more prior lines of treatment and had discontinued ibrutinib due to a lack of efficacy. Most patients (79%) had previously received ibrutinib as a monotherapy. The ORR of 45% and median DOR of 4.3 months were lower than in previous clinical trials of ibrutinib monotherapy for relapsed/refractory MCL; there were also a higher number of prior regimens in the current study [34, 35]. These factors suggest a higher-risk cohort and a potential negative impact on response to subsequent therapy, including lenalidomide. Nonetheless, lenalidomide-based treatment demonstrated meaningful clinical activity in this difficult-to-treat patient population, as demonstrated by a 29% ORR and 14% CR, with a 20-week (95% CI, 2.9 to not available) median DOR. For the DOR analysis, it should be noted that because 82% of responders were censored, the data should be interpreted with caution. With no new safety signals identified, the safety profile in these patients matched the well-established safety shown in multiple studies of lenalidomide monotherapy [22–28].

Prior studies have shown that lenalidomide treatment had significant clinical activity in relapsed/refractory MCL. The MCL-001 EMERGE study reported a 28% ORR (including 8% CR/CR unconfirmed) and 16.6-month DOR with lenalidomide monotherapy in 134 patients with relapsed/refractory MCL after bortezomib treatment. Patients from MCL-001 had received a median of four prior treatment regimens, and 88% had been treated with at least three prior systemic antilymphoma therapies [26]. A UK study reported a 31% ORR, 8% CR, and 22.2-month median DOR with single-agent lenalidomide (6 cycles at 25 mg/day followed by 15 mg/day lower maintenance dose) in 26 patients with relapsed/refractory MCL who had received a median of three prior systemic therapies [25]. The lower DOR of < 5 months in the current study could be a result of ibrutinib resistance. In the randomized MCL-002 (SPRINT) study of 254 patients with relapsed/refractory MCL, the lenalidomide monotherapy group showed higher ORR (40 versus 11%; $P < 0.001$) compared with

Table 5 Documented treatment-emergent all-grade adverse events in $\geq 10\%$ of patients (safety population)

	L (n = 13)		L + R (n = 11)		L + other (n = 34)		Overall (N = 58)	
	No.	%	No.	%	No.	%	No.	%
Hematologic								
Anemia	2	15	3	27	5	15	10	17
Thrombocytopenia	1	8	1	9	7	21	9	16
Neutropenia	1	8	1	9	6	18	8	14
Pancytopenia	1	8	3	27	3	9	7	12
Febrile neutropenia	0	0	0	0	6	18	6	10
Nonhematologic								
Fatigue	4	31	4	36	14	41	22	38
Nausea	2	15	2	18	7	21	11	19
Dizziness	2	15	2	18	7	21	11	19
Dyspnea	2	15	3	27	6	18	11	19
Peripheral edema	0	0	2	18	9	26	11	19
Rash	2	15	1	9	7	21	10	17
Cough	1	8	3	27	7	21	11	19
Decreased appetite	2	15	0	0	5	15	7	12
Diarrhea	0	0	1	9	7	21	8	14
Headache	3	23	1	9	2	6	6	10
Pyrexia	1	8	0	0	5	15	6	10
Vomiting	0	0	2	18	4	12	6	10
Constipation	0	0	0	0	6	18	6	10
Laboratory investigations								
Platelet count decreased	2	15	1	9	3	9	6	10
White blood cell count decreased	1	8	1	9	4	12	6	10

L lenalidomide, L + R lenalidomide plus rituximab

investigator's choice (monotherapy with chlorambucil, cytarabine, gemcitabine, fludarabine, or rituximab), respectively [28]. Median DOR was 16.1 months for lenalidomide and 10.4 months for the investigator's choice group. Lenalidomide in combination with rituximab (R^2) has also shown activity in relapsed/refractory MCL. In a phase I/II dose-finding study, R^2 was well tolerated in MCL, and among 44 patients in phase II, ORR was 57% (CR 36%) and DOR was 18.9 months [30]. A phase II study of iNHL or MCL showed lenalidomide monotherapy followed by R^2 overcame rituximab resistance [29]. In the 14 patients with MCL, ORR after lenalidomide monotherapy and R^2 was 55% for each; DOR to R^2 was 22.1 months. Since responses to lenalidomide in the post-ibrutinib setting are not durable, early referral for allogeneic hematopoietic stem cell transplantation (allo-HCT) should be strongly considered for responding MCL patients without advanced comorbidities [36–38].

There are several limitations to the study, including the retrospective nature of chart review and limited follow-up, which contribute to censoring patients for time-to-event statistics such as DOR. The prevalence of AEs may also be underestimated due to possible under-reporting or other uncontrolled factors such as pre-existing events. Safety summary tables were generated with the expectation of missing data (e.g., grade, treatment-relatedness, seriousness) that might limit the safety analysis. Because of the heterogeneity of regimens combined with lenalidomide, it is difficult to confidently discern the amount of response due to lenalidomide versus the other therapies used in combination, apart from two responses to lenalidomide monotherapy. The two responders to lenalidomide monotherapy represented only 12% of the 17 patients who responded on lenalidomide-containing therapy, further complicating delineation of the effects of lenalidomide with or without other therapies. It would also be beneficial to deduce which patients were previously refractory to rituximab.

As ibrutinib is being used more frequently for patients with MCL, the opportunity now arises to assess the role of other therapies following ibrutinib. Because multiple studies have shown that MCL patients with ibrutinib failure demonstrate poor outcomes with subsequent therapy [16, 17]; it is critical to identify therapies that may provide activity in these patients. Multiple second-generation BTK inhibitors are being investigated to evaluate possible improvements in target specificity, potency, and tolerability through this pathway [39, 40].

Conclusion

Results from this observational study indicate that lenalidomide-based therapy has clinically significant activity as a monotherapy and in combination regimens to treat heavily pretreated patients with refractory or relapsed MCL after ibrutinib therapy or who cannot

tolerate ibrutinib, and thus, lenalidomide addresses an unmet medical need and widens the therapeutic options in a difficult-to-treat patient population.

Additional files

Additional file 1: Table S1. Number of patients per study site. **Table S2.** Lenalidomide combination treatments for L + other group ($n = 34$). **Table S3** Prior systemic anti-lymphoma therapies ($\geq 10\%$ of patients; $N = 58$)*. (DOCX 42 kb)

Acknowledgements

The editorial support was provided by Bio Connections LLC and funded by Celgene Corporation. The authors directed the development of the manuscript and are fully responsible for all content and editorial decisions.

Funding

This study was supported by the Celgene Corporation, Summit, NJ.

Availability of data and materials

The material generated or analyzed during this study are included within the article and its supplementary information files.

Authors' contributions

All authors share equal responsibility for the study and provided equal contribution. All authors read and approved the final manuscript.

Competing interests

MW reports research grants, honoraria, and nonfinancial support from Janssen, Acerta Pharma, Pharmacyclics, and Celgene; research grants and nonfinancial support from Kite Pharma; research grants from Juno Therapeutics, Amgen, Karyopharm Therapeutics, Asana BioSciences, BeiGene, Novartis, Oncoceutics, Oncernal Therapeutics, and Karus; and honoraria from Biolnvent International and Adienne Pharma & Biotech. SJS reports research funding from Celgene during the conduct of the study. IL and TP declare that they have no competing interests. AG reports research funding for the institution from, and was a consultant for, Celgene, Genentech, and Pharmacyclics/J&J and received honoraria from Celgene, Pharmacyclics/J&J, Acerta, and Takeda. SR reports personal fees from Roche, Janssen, AstraZeneca, Celgene, Napp, AbbVie, and MSD. MH was on the speaker's bureau for Celgene and Sanofi, received honorarium from Celgene, received research funding from Takeda and Sanofi, and was a consultant for Janssen R&D. NG reports speaker's bureau and consultancy for Celgene, Gilead, and AbbVie; speaker's bureau, consultancy, and research funding for PCYC and Janssen; consultancy and research funding for SGN; research funding for Genentech; and consultancy and research funding for TG Therapeutics. CBR reports research funding from Celgene, Millennium, and Novartis. MLCB and EB are employees of Celgene Corporation. PM reports personal consulting fees from Celgene and Janssen/Pharmacyclics.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of Lymphoma/Myeloma, University of Texas MD Anderson Cancer Center, 1515 Holcombe Boulevard, Houston, TX, USA. ²Abramson Cancer Center of the University of Pennsylvania, Philadelphia, PA, USA. ³University of Michigan, Ann Arbor, MI, USA. ⁴Sylvester Comprehensive Cancer Center, Division of Hematology Oncology, University of Miami, Miami, FL, USA. ⁵John Theurer Cancer Center at HUMC, Hackensack, NJ, USA. ⁶Department of Haematology, Derriford Hospital and Plymouth University Medical School, Plymouth, UK. ⁷Medical College of Wisconsin & CIMBTR, Milwaukee, WI, USA. ⁸Carolinas HealthCare System, Levine Cancer Institute, Charlotte, NC, USA. ⁹Mayo Clinic Scottsdale/Phoenix, Scottsdale, AZ, USA.

¹⁰Celgene Corporation, Summit, NJ, USA. ¹¹Celgene International Sàrl, Boudry, Switzerland. ¹²Weill Cornell Medical College, New York, NY, USA.

Received: 13 September 2017 Accepted: 17 October 2017
Published online: 02 November 2017

References

- Narurkar R, Alkayem M, Liu D. SOX11 is a biomarker for cyclin D1-negative mantle cell lymphoma. *Biomark Res.* 2016;4:6.
- Swerdlow SH, Campo E, Pileri SA, Harris NL, Stein H, Siebert R, et al. The 2016 revision of the World Health Organization classification of lymphoid neoplasms. *Blood.* 2016;127:2375–90.
- Zhou Y, Wang H, Fang W, Romaguer JE, Zhang Y, Delasalle KB, et al. Incidence trends of mantle cell lymphoma in the United States between 1992 and 2004. *Cancer.* 2008;113:791–8.
- Dreyling M, Geisler C, Hermine O, Kluijn-Nelemans HC, Le Gouill S, Rule S, et al. Newly diagnosed and relapsed mantle cell lymphoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2014; 25 Suppl 3:iii83–iii92.
- Avivi I, Goy A. Refining the mantle cell lymphoma paradigm: impact of novel therapies on current practice. *Clin Cancer Res.* 2015;21:3853–61.
- Cheah CY, Seymour JF, Wang ML. Mantle cell lymphoma. *J Clin Oncol.* 2016; 34:1256–69.
- Rodriguez J, Gutierrez A, Palacios A, Navarrete M, Blancas I, Alarcon J, et al. Rituximab, gemcitabine and oxaliplatin: an effective regimen in patients with refractory and relapsing mantle cell lymphoma. *Leuk Lymphoma.* 2007; 48:2172–8.
- Wang M, Fayad L, Cabanillas F, Hagemester F, McLaughlin P, Rodriguez MA, et al. Phase 2 trial of rituximab plus hyper-CVAD alternating with rituximab plus methotrexate-cytarabine for relapsed or refractory aggressive mantle cell lymphoma. *Cancer.* 2008;113:2734–41.
- Witzig TE, Geyer SM, Kurtin PJ, Colgan JP, Inwards DJ, Micallef IN, et al. Salvage chemotherapy with rituximab DHAP for relapsed non-Hodgkin lymphoma: a phase II trial in the North Central Cancer Treatment Group. *Leuk Lymphoma.* 2008;49:1074–80.
- REVLIMID (lenalidomide) prescribing information. Summit, NJ: Celgene Corporation; 2017.
- Imbruvica (ibrutinib) prescribing information. Sunnyvale, CA: Pharmacyclics, Inc.; 2017.
- Velcade (bortezomib) prescribing information. Cambridge, MA: Millennium Pharmaceuticals, Inc.; 2017.
- Torisel (temsirolimus) prescribing information. Kent, United Kingdom: Pfizer Limited; 2017.
- Imbruvica (ibrutinib) prescribing information. Belgium: Janssen-Cilag International NV; 2017.
- Dreyling M, Jurczak W, Jerkeman M, Silva RS, Rusconi C, Tmemy M, et al. Ibrutinib versus temsirolimus in patients with relapsed or refractory mantle-cell lymphoma: an international, randomised, open-label, phase 3 study. *Lancet.* 2016;387:770–8.
- Cheah CY, Chihara D, Romaguera JE, Fowler NH, Seymour JF, Hagemester FB, et al. Patients with mantle cell lymphoma failing ibrutinib are unlikely to respond to salvage chemotherapy and have poor outcomes. *Ann Oncol.* 2015;26:1175–9.
- Martin P, Maddocks K, Leonard JP, Ruan J, Goy A, Wagner-Johnston N, et al. Postibrutinib outcomes in patients with mantle cell lymphoma. *Blood.* 2016; 127:1559–63.
- Zhang L, Qian Z, Cai Z, Sun L, Wang H, Bartlett JB, et al. Synergistic antitumor effects of lenalidomide and rituximab on mantle cell lymphoma in vitro and in vivo. *Am J Hematol.* 2009;84:553–9.
- Qian Z, Zhang L, Cai Z, Sun L, Wang H, Yi Q, et al. Lenalidomide synergizes with dexamethasone to induce growth arrest and apoptosis of mantle cell lymphoma cells in vitro and in vivo. *Leuk Res.* 2011;35:380–6.
- Song K, Herzog BH, Sheng M, Fu J, McDaniel JM, Chen H, et al. Lenalidomide inhibits lymphangiogenesis in preclinical models of mantle cell lymphoma. *Cancer Res.* 2013;73:7254–64.
- Moros A, Bustany S, Cahu J, Saborit-Villarroya I, Martinez A, Colomer D, et al. Antitumoral activity of lenalidomide in in vitro and in vivo models of mantle cell lymphoma involves the destabilization of cyclin D1/p27KIP1 complexes. *Clin Cancer Res.* 2014;20:393–403.
- Wiernik PH, Lossos IS, Tusciano JM, Justice G, Vose JM, Cole CE, et al. Lenalidomide monotherapy in relapsed or refractory aggressive non-Hodgkin's lymphoma. *J Clin Oncol.* 2008;26:4952–7.
- Habermann TM, Lossos IS, Justice G, Vose JM, Wiernik PH, McBride K, et al. Lenalidomide oral monotherapy produces a high response rate in patients with relapsed or refractory mantle cell lymphoma. *Br J Haematol.* 2009;145: 344–9.
- Witzig TE, Vose JM, Zinzani PL, Reeder CB, Buckstein R, Polikoff JA, et al. An international phase II trial of single-agent lenalidomide for relapsed or refractory aggressive B-cell non-Hodgkin's lymphoma. *Ann Oncol.* 2011;22: 1622–7.
- Eve HE, Carey S, Richardson SJ, Heise CC, Mamidipudi V, Shi T, et al. Single-agent lenalidomide in relapsed/refractory mantle cell lymphoma: results from a UK phase II study suggest activity and possible gender differences. *Br J Haematol.* 2012;159:154–63.
- Goy A, Sinha R, Williams ME, Kalayoglu Besik S, Drach J, Ramchandren R, et al. Single-agent lenalidomide in patients with mantle-cell lymphoma who relapsed or progressed after or were refractory to bortezomib: phase II MCL-001 (EMERGE) study. *J Clin Oncol.* 2013;31:3688–95.
- Zinzani PL, Vose JM, Czuczman MS, Reeder CB, Haioun C, Polikoff J, et al. Long-term follow-up of lenalidomide in relapsed/refractory mantle cell lymphoma: subset analysis of the NHL-003 study. *Ann Oncol.* 2013; 24:2892–7.
- Tmemy M, Lamy T, Walewski J, Belada D, Mayer J, Radford J, et al. Lenalidomide versus investigator's choice in relapsed or refractory mantle cell lymphoma (MCL-002; SPRINT): a phase 2, randomised, multicentre trial. *Lancet Oncol.* 2016;17:319–31.
- Chong EA, Ahmadi T, Aquil NA, Svoboda J, Nasta SD, Mato AR, et al. Combination of lenalidomide and rituximab overcomes rituximab resistance in patients with indolent B-cell and mantle cell lymphomas. *Clin Cancer Res.* 2015;21:1835–42.
- Wang M, Fayad L, Wagner-Bartak N, Zhang L, Hagemester F, Neelapu SS, et al. Lenalidomide in combination with rituximab for patients with relapsed or refractory mantle-cell lymphoma: a phase 1/2 clinical trial. *Lancet Oncol.* 2012;13:716–23.
- Cheson BD, Pfistner B, Juweid ME, Gascoyne RD, Specht L, Horning SJ, et al. Revised response criteria for malignant lymphoma. *J Clin Oncol.* 2007;25: 579–86.
- Cheson BD, Horning SJ, Coiffier B, Shipp MA, Fisher RI, Connors JM, et al. Report of an international workshop to standardize response criteria for non-Hodgkin's lymphomas. NCI Sponsored International Working Group. *J Clin Oncol.* 1999;17:1244.
- Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. *J Am Stat Assoc.* 1958;53:457–81.
- Wang ML, Rule S, Martin P, Goy A, Auer R, Kahl BS, et al. Targeting BTK with ibrutinib in relapsed or refractory mantle-cell lymphoma. *N Engl J Med.* 2013;369:507–16.
- Wang ML, Blum KA, Martin P, Goy A, Auer R, Kahl BS, et al. Long-term follow-up of MCL patients treated with single-agent ibrutinib: updated safety and efficacy results. *Blood.* 2015;126:739–45.
- Epperla N, Hamadani M, Cashen AF, Ahn KW, Oak E, Kanate AS, et al. Predictive factors and outcomes for ibrutinib therapy in relapsed/refractory mantle cell lymphoma—a “real world” study. *Hematol Oncol* 2017;doi: <https://doi.org/10.1002/hon.2380>. Epub ahead of print.
- Fenske TS, Zhang MJ, Carreras J, Ayala E, Burns LJ, Cashen A, et al. Autologous or reduced-intensity conditioning allogeneic hematopoietic cell transplantation for chemotherapy-sensitive mantle-cell lymphoma: analysis of transplantation timing and modality. *J Clin Oncol.* 2014;32:273–81.
- Hamadani M, Saber W, Ahn KW, Carreras J, Cairo MS, Fenske TS, et al. Allogeneic hematopoietic cell transplantation for chemotherapy-unresponsive mantle cell lymphoma: a cohort analysis from the center for international blood and marrow transplant research. *Biol Blood Marrow Transplant.* 2013;19:625–31.
- Wu J, Liu C, Tsui ST, Liu D. Second-generation inhibitors of Bruton tyrosine kinase. *J Hematol Oncol.* 2016;9:80.
- Wu J, Zhang M, Liu D. Acalabrutinib (ACP-196): a selective second-generation BTK inhibitor. *J Hematol Oncol.* 2016;9:21.